







Regio- and Stereoselective Synthesis of (*E*)-1-Bromo-2-iodoalkenes From Internal Alkynes

Masataka Ide, Tetsuo Iwasawa*
iwasawa@rins.ryukoku.ac.jp


Department of Materials Chemistry, Ryukoku University, Seta, Otsu, Shiga 520-2194, Japan



1. Summary a) Ide, M.; Yauchi, Y.; Shioyai, R.; Iwasawa, T. *Tetrahedron* 2014, 70, 8532-8538. b) Japanese Patent No. 2014-153644




4. Halogen sources




X	X'	Yield (%)	Ratio	Recovered SM (%)	Product (X/X')
I	I	26	~ 100:0	60	II
I	Br	76	87:13	19	Brl
I	Cl	24	~ 100:0	32	II
Br	I	81	98:2	0	Brl
Br	Br	-	-	0	-
Br	Cl	-	-	0	-
Cl	I	30	~ 100:0	11	II
Cl	Br	-	-	0	-
Cl	Cl	-	-	0	-

2. Limitation




3. Background

1) Ferreira, 2013 (*ACIE*, vol. 52, pp 7579)



2) Ogilvie, 2006 (*JOC*, vol. 71, pp 3615)



5. Mechanistic aspect

